



Sharp upper bounds on the spectral radius of the signless Laplacian matrix of a graph

A. Dilek (Güngör) Maden^a, Kinkar Ch. Das^{b,*}, A. Sinan Çevik^a

^a Selçuk University, Science Faculty, Department of Mathematics, 42031 Konya, Turkey

^b Department of Mathematics, Sungkyunkwan University, Suwon 440-746, Republic of Korea

ARTICLE INFO

Keywords:

Graph
Spectral radius
Signless Laplacian
Bounds
Degrees
Average degree of neighbors

ABSTRACT

Let $G = (V, E)$ be a simple connected graph. Denote by $D(G)$ the diagonal matrix of its vertex degrees and by $A(G)$ its adjacency matrix. Then the signless Laplacian matrix of G is $Q(G) = D(G) + A(G)$. In this paper, we obtain some new and improved sharp upper bounds on the spectral radius $q_1(G)$ of the signless Laplacian matrix of a graph G .

© 2012 Elsevier Inc. All rights reserved.

1. Introduction

Let $G = (V, E)$ be a simple connected graph with vertex set $V(G) = \{1, 2, \dots, n\}$ and edge set $E(G)$, where $|V(G)| = n$ is the order and $|E(G)| = m$ is the size of G . When i is adjacent to j , we denote this fact by $i \sim j$. For $v_i \in V(G)$, the degree (= number of first neighbors) of the vertex v_i is denoted by d_i . The minimum vertex degree is denoted by δ , the maximum by Δ_1 and the second maximum by Δ_2 . Assuming that the degrees are ordered as $d_1 \geq d_2 \geq \dots \geq d_n$, then $\Delta_1 = d_1$, $\Delta_2 = d_2$ and $\delta = d_n$. The average degree of the neighbors of vertex i is $m_i = \frac{1}{d_i} \sum_{j \sim i} d_j$. The eigenvalues of G are the eigenvalues of the adjacency matrix $A(G)$, given as $\lambda_1(G) \geq \lambda_2(G) \geq \dots \geq \lambda_n(G)$, where, $\lambda_1(G)$ is called the index of G . Consider $D(G)$ as the diagonal matrix of vertex degrees of G . The Laplacian matrix of G is $L(G) = D(G) - A(G)$, its eigenvalues are as displayed as $\mu_1(G) \geq \mu_2(G) \geq \dots \geq \mu_{n-1}(G) \geq \mu_n(G)$ and $\mu_1(G)$ is the spectral radius of Laplacian matrix of graph G . Since $L(G)$ and $A(G)$ are well known, there are many results on their spectra, (see [4,8,10–12,16,20,21,24]).

The matrix $Q(G) = D(G) + A(G)$ was introduced in the classical book of Cvetković, Doob and Sachs on “Spectra of Graphs” [6], but without a name being given to it at that time. Later it was called “quasi-Laplacian matrix” and more recently “signless Laplacian” [3,5,13–15]. Let $q_1(G)$ be the spectral radius of $Q(G)$. Since G is a connected graph then $Q(G)$ is a nonnegative, symmetric and irreducible matrix. Some researchers [25,27] have observed that

$$\mu_1(G) \leq q_1(G) \quad (1)$$

and

$$2\lambda_1(G) \leq q_1(G). \quad (2)$$

These relations immediately imply that any lower bound on $\mu_1(G)$ is a valid lower bound on $q_1(G)$ and that doubling any lower bound on $\lambda_1(G)$ also yields a valid lower bound on $q_1(G)$. A number of upper bounds on $\mu_1(G)$ given as functions of the

* Corresponding author.

E-mail addresses: drdilekgungor@gmail.com (A. Dilek (Güngör) Maden), kinkardas2003@googlemail.com (K.Ch. Das), sinan.cevik@selcuk.edu.tr (A. Sinan Çevik).

degree and of the average degree of the neighbors of a vertex have been proposed in the literature. In [2,23], it was gathered some of them and these bounds are

$$\mu_1(G) \leq \max_{i \in V} \{2d_i\}, \quad [23] \quad (3)$$

$$\mu_1(G) \leq \max_{i \in V} \{d_i + m_i\}, \quad [22], \quad (4)$$

$$\mu_1(G) \leq \max_{i \in V} \left\{ d_i + \sqrt{d_i m_i} \right\}, \quad [23, 29], \quad (5)$$

$$\mu_1(G) \leq \max_{i \in V} \left\{ \sqrt{2d_i(d_i + m_i)} \right\}, \quad [19], \quad (6)$$

$$\mu_1(G) \leq \max_{i \in V} \left\{ \frac{d_i + \sqrt{d_i^2 + 8d_i m_i}}{2} \right\}, \quad [17, 23], \quad (7)$$

$$\mu_1(G) \leq \max_{i \sim j} \{d_i + d_j\}, \quad [1], \quad (8)$$

$$\text{and } \mu_1(G) \leq \max_{i \sim j} \left\{ \frac{d_i + d_j + \sqrt{(d_i - d_j)^2 + 4m_i m_j}}{2} \right\}, [9, 28]. \quad (9)$$

In the above bounds (3),(4),(5) and (7),(8),(9) are valid for $q_1(G)$ (see [2,3,11,23,26]). In this paper, we obtain some new and improved sharp upper bounds on it.

2. Sharp upper bounds for the spectral radius of the signless Laplacian matrix of a graph

We first list some known results which will be used in this paper.

Lemma 1 [18]. Let M be irreducible non-negative matrix. Then $\rho(M)$ is an eigenvalue of M and there is a positive vector X such that $MX = \rho(M)X$.

Lemma 2 [5]. Let $M = (m_{ij})$ be an $n \times n$ non-negative matrix and let $R_i(M)$ be the i th row sum of M , i.e., $R_i(M) = \sum_{j=1}^n m_{ij} (1 \leq i \leq n)$. Then

$$\min \{R_i(M) : 1 \leq i \leq n\} \leq \rho(M) \leq \max \{R_i(M) : 1 \leq i \leq n\}. \quad (10)$$

If M is irreducible, then each equality holds if and only if $R_1 = R_2 = \dots = R_n$.

Theorem 3 [30]. Let $A = (a_{ij})$ be an $n \times n$ irreducible non-negative matrix. Then

$$\min \left\{ \sqrt{\frac{\sum_{j=1}^n a_{ij} M_j}{R_i}}, 1 \leq i \leq n \right\} \leq \rho(A) \leq \max \left\{ \sqrt{\frac{\sum_{j=1}^n a_{ij} M_j}{R_i}}, 1 \leq i \leq n \right\}, \quad (11)$$

where $R_i = \sum_{j=1}^n a_{ij}$, $M_i = \sum_{j=1}^n a_{ij} R_j$, $M'_i = \frac{M_i}{R_i}$ and $\rho(A)$ denotes the spectral radius of A . Moreover, if A^2 is irreducible, then any equality holds in (11) if and only if $M'_1 = M'_2 = \dots = M'_n$; and if A^2 is reducible, then any equality holds in (11) if and only if there exist the permutation matrix P such that

$$PAP^T = \begin{pmatrix} 0_r & A_1 \\ A_2 & 0_{n-r} \end{pmatrix},$$

and $M'_{\sigma(1)} = \dots = M'_{\sigma(r)}$, $M'_{\sigma(r+1)} = \dots = M'_{\sigma(n)}$, where σ is a permutation on the set $\{1, 2, \dots, n\}$ which corresponds to the permutation matrix P .

Corollary 4 [30]. Let $A = (a_{ij})$ be an $n \times n$ irreducible non-negative matrix. Then

$$\min \{M'_i : 1 \leq i \leq n\} \leq \rho(A) \leq \max \{M'_i : 1 \leq i \leq n\}, \quad (12)$$

where M'_i be as in Theorem 3. Then equality holds in (12) if and only if $M'_1 = M'_2 = \dots = M'_n$.

Corollary 5 [30]. Let $A = (a_{ij})$ be an $n \times n$ irreducible non-negative matrix. Then

$$\min \left\{ \sqrt{M_i} : 1 \leq i \leq n \right\} \leq \rho(A) \leq \max \left\{ \sqrt{M_i} : 1 \leq i \leq n \right\}, \quad (13)$$

where M_i be as in Theorem 3. Moreover, if A^2 is irreducible, then any equality holds in (13) if and only if $R_1 = R_2 = \dots = R_n$; and if A^2 is reducible, then any equality holds in (13) if and only if there exist the permutation matrix P such that

$$PAP^T = \begin{pmatrix} O_r & A_1 \\ A_2 & O_{n-r} \end{pmatrix},$$

and $R_{\sigma(1)} = \dots = R_{\sigma(r)}$, $R_{\sigma(r+1)} = \dots = R_{\sigma(n)}$, where σ is a permutation on the set $\{1, 2, \dots, n\}$ which corresponds to the permutation matrix P .

Now we give our main results. Throughout this paper, G will denote a simple connected graph on n vertices unless stated otherwise.

Theorem 6. Let $b_i \in \mathbb{R}^+$, $1 \leq i \leq n$. Also let $b'_i = \frac{1}{b_i} \sum_{j \sim i} b_j$, $c_i = b_i(d_i + b'_i)$, $c'_i = \frac{\sum_{j \sim i} c_j}{c_i}$ and $k_i = d_i + c'_i$. Then

(a)

$$q_1(G) \leq \max_{1 \leq i \leq n} \left\{ \sqrt{d_i k_i + \frac{\sum_{j \sim i} (d_j c_j + \sum_{k \sim j} c_k)}{c_i}} \right\}. \quad (14)$$

If Q^2 is irreducible, then equality holds in (14) if and only if $k_1 = k_2 = \dots = k_n$ and if Q^2 is reducible, then equality holds in (14) if and only if there exists the permutation matrix P such that

$$PQ(G)P^T = \begin{pmatrix} O_r & Q_1 \\ Q_2 & O_{n-r} \end{pmatrix},$$

with $k_{\sigma(1)} = k_{\sigma(2)} = \dots = k_{\sigma(r)}$ and $k_{\sigma(r+1)} = k_{\sigma(r+2)} = \dots = k_{\sigma(n)}$, where σ is a permutation on the set $\{1, 2, \dots, n\}$ which corresponds to the permutation matrix P .

(b)

$$q_1(G) \leq \max_{i \sim j} \left\{ \frac{d_i + d_j + \sqrt{(d_i - d_j)^2 + 4b'_i b'_j}}{2} \right\}, \quad (15)$$

with equality holds in (15) if and only if G is either a regular graph or a bipartite semi-regular graph.

(c)

$$q_1(G) \leq \max_{1 \leq i \leq n} \{k_i\}, \quad (16)$$

with equality holds in (16) if and only if $k_1 = k_2 = \dots = k_n$.

(d)

$$q_1(G) \leq \max_{1 \leq i \leq n} \left\{ \sqrt{\frac{c_i(d_i + c'_i)}{b_i}} \right\}. \quad (17)$$

If Q^2 is irreducible, then the equality holds in (17) if and only if $d_1 + b'_1 = d_2 + b'_2 = \dots = d_n + b'_n$; and if Q^2 is reducible, then the equality holds in (17) if and only if there exists the permutation matrix P such that

$$PQ(G)P^T = \begin{pmatrix} O_r & Q_1 \\ Q_2 & O_{n-r} \end{pmatrix},$$

with $d_{\sigma(1)} + b'_{\sigma(1)} = d_{\sigma(2)} + b'_{\sigma(2)} = \dots = d_{\sigma(r)} + b'_{\sigma(r)}$ and $d_{\sigma(r+1)} + b'_{\sigma(r+1)} = d_{\sigma(r+2)} + b'_{\sigma(r+2)} = \dots = d_{\sigma(n)} + b'_{\sigma(n)}$, where σ is a permutation on the set $\{1, 2, \dots, n\}$ which corresponds to the permutation matrix P .

Proof. Let $B = \text{diag}(b_1, b_2, \dots, b_n)$ be an $n \times n$ diagonal matrix. Consider the matrix $B^{-1}Q(G)B$. Now the (i, j) th element of $B^{-1}Q(G)B$ is

$$\begin{cases} d_i & \text{if } i = j, \\ \frac{b_j}{b_i} & \text{if } i \sim j, \\ 0 & \text{otherwise.} \end{cases}$$

It is easy to see that the inequality (14) holds from Theorem 3 as

$$R_i = \sum_{j=1}^n a_{ij} = d_i + \frac{1}{b_i} \sum_{j \sim i} b_j = \frac{c_i}{b_i},$$

$$M_i = d_i R_i + \sum_{j \sim i} \frac{b_j}{b_i} R_j = \frac{1}{b_i} \left(d_i c_i + \sum_{j \sim i} c_j \right) = \frac{k_i c_i}{b_i}$$

and

$$\sum_{j=1}^n a_{ij} M_j = d_i M_i + \sum_{j \sim i} a_{ij} M_j = \frac{d_i k_i c_i}{b_i} + \sum_{j \sim i} \frac{k_j c_j}{b_i}.$$

Moreover, the proof of the second part of (a) is directly follows from [Theorem 3](#).

Next, we prove that (15) holds. From [Lemma 1](#), there exists a positive eigenvector $\mathbf{X} = (x_1, x_2, \dots, x_n)^T$ of $B^{-1}Q(G)B$ corresponding to $q_1(B^{-1}Q(G)B)$. We can assume that one of the eigencomponents, say x_i , is equal to 1 and the other eigencomponents are less than or equal to 1, i.e., $x_i = 1$ and $x_k \leq 1$, $1 \leq k \leq n$. Also, let $x_j = \max_k \{x_k : k \sim i\}$. From

$$(B^{-1}Q(G)B)X = q_1(G)X, \quad (18)$$

we have

$$q_1(G) = d_i + \frac{1}{b_i} \sum_{k:k \sim i} b_k x_k \leq d_i + \frac{1}{b_i} \sum_{k:k \sim i} b_k x_j \quad (19)$$

and

$$q_1(G)x_j = d_j x_j + \frac{1}{b_j} \sum_{k:k \sim j} b_k x_k \leq d_j x_j + \frac{1}{b_j} \sum_{k:k \sim j} b_k. \quad (20)$$

From (19) and (20), we have

$$q_1(G)^2 - (d_i + d_j)q_1(G) + d_i d_j - \left(\frac{1}{b_i} \sum_{k:k \sim i} b_k \right) \left(\frac{1}{b_j} \sum_{k:k \sim j} b_k \right) \leq 0.$$

Thus we have

$$q_1(G) \leq \frac{d_i + d_j + \sqrt{(d_i - d_j)^2 + 4b'_i b'_j}}{2}.$$

Hence (15) holds.

Moreover, one can see easily that the equality holds in (15) for regular graph or for bipartite semiregular graph. (16) and (17) with equality follow from [Corollaries 4 and 5](#), respectively. The proof is complete. \square

Remark 7. From [Theorem 6](#), we have the following known results.

1. Taking $b_i = 1$ in (15), we have upper bound (8).
2. Taking $b_i = d_i$ in (15), we have upper bound (9).
3. Taking $b_i = 1$ in (16), we have upper bound (4).
4. Taking $b_i = 1$ in (17), we have upper bound (6).

In particular, from [Theorem 6](#), we have the following results.

Corollary 8. For a graph G , as depicted previously, the following inequalities hold:

$$q_1(G) \leq \max_i \left\{ \sqrt{d_i(d_i + m_i) + s_i} \right\}, \quad (21)$$

$$q_1(G) \leq \max_i \left\{ \sqrt{d_i \left(d_i + \frac{s_i}{d_i + m_i} \right) + \frac{\sum_{j \sim i} d_j [d_j(d_j + m_j) + s_j]}{d_i(d_i + m_i)}} \right\} \quad (22)$$

and

$$q_1(G) \leq \max_i \left\{ d_i + \frac{s_i}{d_i + m_i} \right\}, \quad (23)$$

where $s_i = \sum_{j \sim i} \frac{d_j(d_j + m_j)}{d_i}$.

Proof. Taking $b_i = 1$ and $b_i = d_i$ in (14) and $b_i = d_i$ in (16), respectively, we have the required results. \square

Remark 9. Since Corollary 8 is a consequence of Theorem 6, it is easy to conclude that the equality conditions in (14) and (16) are also hold.

Lemma 10 [7]. Let G be a connected graph. Then $d_1 + m_1 = d_2 + m_2 = \dots = d_n + m_n$ if and only if G is a regular graph or G is a regular bipartite graph.

Theorem 11. Let G be a connected graph. Then

$$q_1(G) \leq \max_{1 \leq i \leq n} \{d_i + b'_i\} \quad (24)$$

and

$$q_1(G) \leq \max_i \left\{ \frac{d_i + \sqrt{d_i^2 + \frac{4c_i c'_i}{b_i}}}{2} \right\}, \quad (25)$$

where $b_i \in \mathbb{R}^+$, $b'_i = \frac{1}{b_i} \sum_{j \sim i} b_j$, $c_i = b_i(d_i + b'_i)$, $c'_i = \sum_{j \sim i} c_j$. Moreover, both the equality hold if and only if $d_1 + b'_1 = d_2 + b'_2 = \dots = d_n + b'_n$.

Proof. Let $X = (x_1, x_2, \dots, x_n)^T$ be an eigenvector corresponding to the eigenvalue $q_1(G)$ of $B^{-1}Q(G)B$. We assume that one eigencomponent x_i is equal to 1 and the other eigencomponents are less than or equal to 1, that is, $x_i = 1$ and $0 < x_k \leq 1$, for all k .

From the i th equation of (18), we have

$$q_1(G)x_i = d_i x_i + \sum_{j \sim i} \frac{b_j x_j}{b_i},$$

i.e., $q_1(G) = d_i + \sum_{j \sim i} \frac{b_j x_j}{b_i}$. (26)

From above the first bound follows. Moreover, the equality holds in (24) if and only if $d_i + b'_i$ ($1 \leq i \leq n$) is a constant. Again from the j th equation of (18),

$$q_1(G)x_j = d_j x_j + \sum_{k \sim j} \frac{b_k x_k}{b_j}.$$

Multiplying both sides of (26) by $q_1(G)$ and substituting this value $q_1(G)x_j$, we get

$$\begin{aligned} q_1^2(G) &= d_i q_1(G) + \sum_{j \sim i} \left\{ \frac{b_j}{b_i} \left[d_j x_j + \sum_{k \sim j} \frac{b_k x_k}{b_j} \right] \right\} \\ &= d_i q_1(G) + \sum_{j \sim i} \frac{b_j d_j}{b_i} x_j + \frac{1}{b_i} \sum_{j \sim i} \sum_{k \sim j} b_k x_k \leq d_i q_1(G) + \sum_{j \sim i} \frac{b_j d_j}{b_i} + \sum_{j \sim i} \frac{b_j b'_j}{b_i} \text{ as } x_j, x_k \leq 1 \\ &= d_i q_1(G) + \sum_{j \sim i} \frac{b_j (d_j + b'_j)}{b_i} = d_i q_1(G) + \frac{c_i c'_i}{b_i} \end{aligned} \quad (27)$$

from above the second bound follows.

Now suppose that the equality holds in (25). Then all inequalities in the above argument must be equalities. From equality in (27), we get $x_j = 1$ for all j such that $j \sim i$ and $x_k = 1$ for all k such that $k \sim j$ and $j \sim i$. From this one can easily show that $x_i = 1$ for all $i \in V$. Thus we have $d_1 + b'_1 = d_2 + b'_2 = \dots = d_n + b'_n$.

Conversely, one can easily see that the equality holds in (25) for $d_1 + b'_1 = d_2 + b'_2 = \dots = d_n + b'_n$. \square

Corollary 12. Let G be a connected graph. Then

$$q_1(G) \leq \max_{1 \leq i \leq n} \left\{ \frac{d_i + \sqrt{d_i^2 + \frac{8d_i m_i b_i}{b_i}}}{2} \right\}, \quad (28)$$

where $b'_i = \frac{1}{b_i} \sum_{j \sim i} b_j$, $b_\ell = \max_{1 \leq i \leq n} b_i$ and $m_i = \frac{1}{d_i} \sum_{j \sim i} d_j$.

Corollary 13. Let G be a connected graph. Then

$$q_1(G) \leq \max_{1 \leq i \leq n} \left\{ \frac{d_i + \sqrt{d_i^2 + \frac{4}{d_i} \sum_{j \sim i} d_j (d_j + m_j)}}{2} \right\}.$$

The equality holds if and only if G is a regular graph or G is a regular bipartite graph.

Proof. Taking $b_i = d_i$ in (25), the above bound follows. By Lemma 10 and Theorem 11, the above equality holds if and only if G is a regular graph or G is a regular bipartite graph. \square

Corollary 14. Let G be a connected graph. Then

$$q_1(G) \leq \max_{1 \leq i \leq n} \left\{ d_i + \frac{1}{\sqrt{d_i}} \sum_{j \sim i} \sqrt{d_j} \right\}, \quad (29)$$

with equality if and only if $d_i + \frac{1}{\sqrt{d_i}} \sum_{j \sim i} \sqrt{d_j}$ ($1 \leq i \leq n$) is a constant.

Proof. Taking $b_i = \sqrt{d_i}$ in (24), the result follows. \square

Remark 15. From the Cauchy–Schwarz inequality, it is easy to see that bound (29) is always better than bound (5).

Remark 16. From Theorem 11 and Corollary 12, respectively, we have the following known results:

1. Taking $b_i = 1$ and $b_i = d_i$ in (24), we have upper bounds (3) and (4).
2. Taking $b_i = 1$ in (25) and (28), we have upper bound (7).

Let Γ be the class of graphs $H = (V, E)$ such that H is connected graph with $V(H) = \{1\} \cup V_1 \cup V_2$,

$$d_1 = \Delta_1, \quad V_1 = \{k \in N_1 : d_k = \delta\}, \quad V_2 = \{k \notin N_1 : d_k = \Delta_2\}$$

and

$$(\Delta_2 - \delta)(2\Delta_2 - \Delta_1) = \Delta_1 - \Delta_2.$$

The spectral radius of the signless Laplacian matrix of $H \in \Gamma$ is given by:

$$q_1(H) = 2\Delta_2 = \frac{\Delta_1 + 2\delta - 1 + \sqrt{(\Delta_1 - 2\delta + 1)^2 + 4\Delta_1}}{2}.$$

Denote by H_{n-1, Δ_2} , a connected graph with maximum degree $n-1$ and the second maximum degree Δ_2 such that $\Delta_2 = \delta < n-1$ (δ is minimum vertex degree). Thus H_{n-1, Δ_2} is a $(n-1, \Delta_2)$ -semiregular graph with $\Delta_2 < n-1$. One can see easily that the spectral radius of the signless Laplacian matrix of H_{n-1, Δ_2} is given by the following equation

$$q_1^2 - (n + 2\Delta_2 - 2)q_1 + 2(n-1)(\Delta_2 - 1) = 0.$$

Thus the spectral radius of the signless Laplacian matrix of H_{n-1, Δ_2} is given by

$$q_1 = \frac{n + 2\Delta_2 - 2 + \sqrt{(n - 2\Delta_2)^2 + 4(n-1)}}{2}.$$

We now give another upper bound on the spectral radius of the signless Laplacian matrix of a graph.

Theorem 17. Let G be a connected graph with maximum degree Δ_1 and second maximum degree Δ_2 . Then

$$q_1 \leq \max_{i \sim j} \left\{ \frac{d_i + 2d_j - 1 + \sqrt{(d_i - 2d_j + 1)^2 + 4d_i}}{2} \right\}, \quad (30)$$

with equality holds in (30) if and only if G is isomorphic to a regular graph or $G \cong H_{n-1, \Delta_2}$ or $G \in \Gamma$.

Proof. Let $\mathbf{X} = (x_1, x_2, \dots, x_n)^T$ be an eigenvector of $Q(G)$ corresponding to an eigenvalue q_1 . We can assume that one eigencomponent x_i is equal to 1 and the other eigencomponents are less than or equal to 1, that is, $x_i = 1$ and $0 < x_k \leq 1$, for all k . We have

$$Q(G)\mathbf{X} = q_1\mathbf{X}. \quad (31)$$

Let $x_j = \max_{k:k \neq i} x_k$. From the i th equation of (31),

$$q_1 x_i = d_i x_i + \sum_{k:k \sim i} x_k, \quad \text{i.e., } q_1 \leq d_i + d_i x_j. \quad (32)$$

From the j th equation of (31),

$$\begin{aligned} q_1 x_j &= d_j x_j + \sum_{k:k \sim j} x_k, \\ \text{i.e., } q_1 x_j &\leq d_j x_j + 1 + (d_j - 1)x_j, \\ \text{i.e., } (q_1 - 2d_j + 1)x_j &\leq 1. \end{aligned} \quad (33)$$

From (32) and (33), we get

$$(q_1 - d_i)(q_1 - 2d_j + 1) \leq d_i,$$

i.e.,

$$q_1^2 - (d_i + 2d_j - 1)q_1 + 2d_i(d_j - 1) \leq 0,$$

i.e.,

$$q_1 \leq \frac{d_i + 2d_j - 1 + \sqrt{(d_i - 2d_j + 1)^2 + 4d_i}}{2}.$$

The first part of the proof is over.

Now suppose that equality holds in (30). Then all inequalities in the above argument must be equalities. In particular, from (32) we get

$$x_k = x_j \text{ for all } k, k \sim i.$$

Also from (33) we get

$$x_k = x_j \text{ for all } k, \quad k \sim j, \quad k \neq i \text{ and } i \sim j.$$

Let $V_1 = \{k : x_k = x_j\}$. If $V_1 \neq V(G) \setminus \{i\}$, then there exist vertices $p \in V_1$, $q \notin V_1$, $q \neq i$ such that $p \sim q$ as G is connected. Thus we have $x_q < x_j$ as x_j is the second maximum eigencomponent. For vertex $p \in V(G)$, from above, we must have $x_q = x_j$, a contradiction. Thus $V_1 = V(G) \setminus \{i\}$. If $x_i = 1$, then

$$q_1 = 2d_i, \quad i = 1, 2, \dots, n.$$

Hence G is a regular graph.

Otherwise, $x_j < 1$. Now we consider two cases (i) $d_i = n - 1$, (ii) $d_i < n - 1$.

Case (i) : $d_i = n - 1$. In this case vertex i is adjacent to all the remaining vertices in G . One can see easily that any two vertices j and k in $V(G) \setminus \{i\}$, $d_j = d_k < n - 1$. Thus we have $\Delta_2 = d_2 = d_3 = \dots = d_n = \delta$. Hence $G \cong H_{n-1, \Delta_2}$.

Case (ii) : $d_i < n - 1$. In this case $V_1 = \{k \in N_i\}$ and $V_2 = V(G) \setminus (V_1 \cup \{i\})$. One can see easily that any two vertices j and k in V_1 , $d_j = d_k$ and also we have $d_r = d_s$, for $r, s \in V_2$. Moreover, we have

$$q_1 = d_1(1 + x_j), \quad (34)$$

$$q_1 = 2d_j - 1 + \frac{1}{x_j} \quad j \in V_1, \quad (35)$$

$$\text{and } q_1 = 2d_k \quad k \in V_2. \quad (36)$$

From (34) and (36), we get $d_1 > d_k$, $k \in V_2$. From (35) and (36), we get $d_k > d_j$, $k \in V_2$, $j \in V_1$. Thus we have $d_1 = \Delta_1$, $d_k = \Delta_2$ and $d_j = \delta$, $k \in V_2$ and $j \in V_1$. Moreover, $(\Delta_2 - \delta)(2\Delta_2 - \Delta_1) = \Delta_1 - \Delta_2$, by (34)–(36).

Hence $G \in \Gamma$.

Conversely, one can see easily that the equality holds in (30) for regular graph or for H_{n-1, Δ_2} or for $G, G \in \Gamma$. \square

Acknowledgements

The first and third authors are partially supported by TUBITAK and the Scientific Research Office of Selcuk University (BAP). The second author is partially supported by the Faculty research Fund, Sungkyunkwan University, 2012 and Sungkyunkwan University BK21 Project, BK21 Math Modeling HRD Div. Sungkyunkwan University, Suwon, Republic of Korea.

References

- [1] W.N. Anderson, T.D. Morley, Eigenvalues of the Laplacian of a graph, *Linear Multilinear Algebra* 18 (1985) 141–145.
- [2] V. Brankov, P. Hansen, D. Stevanović, Automated conjectures on upper bounds for the largest Laplacian eigenvalue of graphs, *Linear Algebra Appl.* 414 (2006) 407–424.
- [3] D. Cvetković, P. Rowlinson, S. Simić, Signless Laplacian of finite graphs, *Linear Algebra Appl.* 423 (2007) 155–171.
- [4] A. Berman, X.-D. Zhang, Lower bounds for the eigenvalues of Laplacian matrices, *Linear Algebra Appl.* 316 (2000) 13–20.
- [5] D. Cvetković, Signless Laplacians and line graphs, *Bulletin T. CXXXI de l'Académie serbe des sciences et des arts* (2005) Classe des Sciences mathématiques et naturelles Sciences mathématiques 30 (2005) 85–92.
- [6] D. Cvetković, M. Doob, H. Sachs, *Spectra of Graphs*, Academic Press, New York, 1980 (second revised ed., Barth, Heidelberg, 1995).
- [7] K.C. Das, Maximizing the sum of the squares of the degrees of a graph, *Discrete Math.* 285 (2004) 57–66.
- [8] K.C. Das, An improved upper bound for Laplacian graph eigenvalues, *Linear Algebra Appl.* 368 (2003) 269–278.
- [9] K.C. Das, A characterization on graphs which achieve the upper bound for the largest Laplacian eigenvalue of graphs, *Linear Algebra Appl.* 368 (2003) 269–278.
- [10] K.C. Das, The largest two Laplacian eigenvalues of a graph, *Linear Multilinear Algebra* 52 (6) (2004) 441–460.
- [11] K.C. Das, The Laplacian spectrum of a graph, *Comput. Math. Appl.* 48 (2004) 715–724.
- [12] K.C. Das, Sharp lower bounds on the Laplacian eigenvalues of trees, *Linear Algebra Appl.* 384 (2004) 155–169.
- [13] K.C. Das, On conjectures involving second largest signless Laplacian eigenvalue of graphs, *Linear Algebra Appl.* 432 (2010) 3018–3029.
- [14] K.C. Das, Proof of conjecture involving the second largest signless Laplacian eigenvalue and the index of graphs, *Linear Algebra Appl.* 435 (2011) 2420–2424.
- [15] K.C. Das, Proofs of conjectures involving the largest and the smallest signless Laplacian eigenvalues of graphs, *Discrete Math.* 312 (2012) 992–998.
- [16] R. Grone, R. Merris, V.S. Sunder, The Laplacian spectral of graphs, *SIAM J. MAA* 11 (1990) 218–239.
- [17] J.M. Guo, A new upper bound for the Laplacian spectral radius of graphs, *Linear Algebra Appl.* 400 (2005) 61–66.
- [18] R.A. Horn, C.R. Johnson, *Matrix Analysis*, Cambridge University Press, New York, 1985.
- [19] J.-S. Li, Y.P. Pan, De Caen's inequality and bounds on the largest Laplacian eigenvalue of a graph, *Linear Algebra Appl.* 328 (2001) 153–160.
- [20] J.-S. Li, X.D. Zhang, On the Laplacian eigenvalues of a graph, *Linear Algebra Appl.* 285 (1998) 305–307.
- [21] R. Merris, Laplacian matrices of graphs: a survey, *Linear Algebra Appl.* 197 (198) (1994) 143–176.
- [22] R. Merris, A note on Laplacian graph eigenvalues, *Linear Algebra Appl.* 285 (1988) 33–35.
- [23] C.S. Oliveira, L.S. Lima, N.M. Abreu, P. Hansen, Bounds on the index of the signless Laplacian of a graph, *Discrete Appl. Math.* 158 (2010) 355–360.
- [24] O. Rojo, R. Soto, H. Rojo, An always nontrivial upper bound for graph eigenvalues, *Linear Algebra Appl.* 312 (2000) 155–159.
- [25] J.L. Shu, Y. Hong, R.K. Wen, A sharp upper bound on the largest eigenvalue of the Laplacian matrix of a graph, *Linear Algebra Appl.* 347 (2002) 123–129.
- [26] S. Tan, J. Guo, J. Qi, The spectral radius of Laplacian matrices and quasi-Laplacian matrices of graphs, *J. Eng. Math. (China)* 20 (6) (2003) 69–74.
- [27] C. Yan, Properties of spectra of graphs and line graphs, *Appl. Math. J. Chin. Uni. Ser. B* 3 (2002) 371–376.
- [28] X.-D. Zhang, The Laplacian eigenvalues of mixed graphs, *Linear Algebra Appl.* 362 (2003) 109–119.
- [29] X.-D. Zhang, Two sharp upper bounds for Laplacian eigenvalues, *Linear Algebra Appl.* 376 (2004) 207–213.
- [30] X.-D. Zhang, J.S. Li, Spectral radius of non-negative matrices and digraphs, *Acta Math. Sin.* 18 (2002) 293–300.