

## SPECTRAL RADIUS AND AVERAGE 2-DEGREE SEQUENCE OF A GRAPH

YU-PEI HUANG

*Department of Applied Mathematics  
I-Shou University  
Kaohsiung City 84001, Taiwan  
pei.am91g@nctu.edu.tw*

CHIH-WEN WENG

*Department of Applied Mathematics  
National Chiao Tung University  
Hsinchu City 30010, Taiwan  
weng@math.nctu.edu.tw*

Received 12 January 2013

Revised 17 December 2013

Accepted 28 January 2014

Published 24 February 2014

In a simple connected graph, the average 2-degree of a vertex is the average degree of its neighbors. With the average 2-degree sequence and the maximum degree ratio of adjacent vertices, we present a sharp upper bound of the spectral radius of the adjacency matrix of a graph, which improves a result in [Y. H. Chen, R. Y. Pan and X. D. Zhang, Two sharp upper bounds for the signless Laplacian spectral radius of graphs, *Discrete Math. Algorithms Appl.* **3**(2) (2011) 185–191].

*Keywords:* Graph; adjacency matrix; spectral radius; average 2-degree.

Mathematics Subject Classification: 05C50

### 1. Introduction

Let  $G = (V, E)$  be a simple connected graph with vertex set  $V(G)$ , edge set  $E(G)$  and  $n = |V(G)|$ . For any vertex  $v \in V(G)$ , let  $d_v$  denote the degree of  $v$ , define the *average 2-degree*  $M_v$  of  $v$  to be the average degree of the neighbors of  $v$ . In other words,  $M_v = \sum_{u \sim v} d_u/d_v$ , where  $u \sim v$  means vertices  $u$  and  $v$  are adjacent. Throughout the paper, label the vertices of  $G$  by  $1, 2, \dots, n$  such that  $M_1 \geq M_2 \geq \dots \geq M_n$ . Let  $A = (a_{ij})$  be the *adjacency matrix* of  $G$ , a binary matrix of order  $n$  such that for any pair  $i, j \in V(G)$ ,  $a_{ij} = 1$  if and only if  $i, j$  are adjacent in  $G$ . The *spectral radius*  $\rho(G)$  of  $G$  is the largest eigenvalue of its adjacency matrix; this parameter has been studied by many authors [1, 2, 5–9, 11–13] and can be

used to induce some other bounds such as the upper bounds of signless Laplacian eigenvalues [3, 4].

The following theorem is well-known and referred as Perron–Frobenius Theorem [10, Chap. 2].

**Theorem 1.1.** *If  $B$  is a nonnegative irreducible  $n \times n$  matrix with largest eigenvalue  $\rho(B)$  and row-sums  $r_1, r_2, \dots, r_n$ , then*

$$\rho(B) \leq \max_{1 \leq i \leq n} r_i$$

*with equality if and only if the row-sums of  $B$  are all equal.*

In this paper, we pay attention to the upper bounds of spectral radius of graphs in terms of the average 2-degree sequence. By setting  $B = U^{-1}AU$ , where  $U = \text{diag}(d_1, d_2, \dots, d_n)$ , the following fact is easily seen from Theorem 1.1.

**Theorem 1.2.**

$$\rho(G) \leq M_1$$

*with equality if and only if  $M_1 = M_2 = \dots = M_n$ .*

A graph for which equality holds in Theorem 1.2 is called *pseudo-regular* in [13].

In 2011 [3, Theorem 2.1], Chen, Pan and Zhang gave the following bound.

**Theorem 1.3.** *Let  $a := \max\{d_i/d_j \mid 1 \leq i, j \leq n\}$ . Then*

$$\rho(G) \leq \frac{M_2 - a + \sqrt{(M_2 + a)^2 + 4a(M_1 - M_2)}}{2},$$

*with equality if and only if  $G$  is pseudo-regular.*

We will show in Corollary 3.3 that Theorem 1.3 is indeed a generalization of Theorem 1.2. Moreover, we give the following theorem to generalize Theorem 1.3.

**Theorem 1.4.** *For any  $b \geq \max\{d_i/d_j \mid i \sim j\}$  and  $1 \leq \ell \leq n$ ,*

$$\rho(G) \leq \frac{M_\ell - b + \sqrt{(M_\ell + b)^2 + 4b \sum_{i=1}^{\ell-1} (M_i - M_\ell)}}{2},$$

*with equality if and only if  $G$  is pseudo-regular.*

Note that Theorem 1.3 is a special case of Theorem 1.4 by taking  $b = a$  and  $\ell = 2$ . The proof of Theorem 1.4 is a subtle application of Perron–Frobenius Theorem. This idea was previously employed in [9, 11]. Indeed, our proof is an edited version of the proof of [9, Theorem 1.7].

We provide some examples of pseudo-regular graphs that are not regular in Example 2.1. The lowest upper bound among the choices of  $b$  and  $\ell$  is investigated in Sec. 3.

## 2. Proof of Theorem 1.4

**Proof of Theorem 1.4.** For each  $1 \leq i \leq \ell - 1$ , let  $x_i \geq 1$  be a variable to be determined later. Let  $U = \text{diag}(d_1 x_1, \dots, d_{\ell-1} x_{\ell-1}, d_\ell, \dots, d_n)$  be a diagonal matrix of size  $n \times n$ . Consider the matrix  $B = U^{-1} A U$ . Note that  $A$  and  $B$  have the same eigenvalues. Let  $r_1, r_2, \dots, r_n$  be the row-sums of  $B$ . Then for  $1 \leq i \leq \ell - 1$  we have

$$\begin{aligned} r_i &= \sum_{k=1}^{\ell-1} \frac{1}{d_i x_i} a_{ik} d_k x_k + \sum_{k=\ell}^n \frac{1}{d_i x_i} a_{ik} d_k \\ &= \frac{1}{x_i} \sum_{k=1}^{\ell-1} (x_k - 1) a_{ik} \frac{d_k}{d_i} + \frac{1}{x_i} \sum_{k=1}^n a_{ik} \frac{d_k}{d_i} \\ &\leq \frac{b}{x_i} \left( \sum_{k=1, k \neq i}^{\ell-1} x_k - (\ell - 2) \right) + \frac{1}{x_i} M_i, \end{aligned} \quad (2.1)$$

since  $a_{ik} d_k / d_i \leq b$ . Similarly for  $\ell \leq j \leq n$  we have

$$\begin{aligned} r_j &= \sum_{k=1}^{\ell-1} x_k a_{jk} \frac{d_k}{d_j} + \sum_{k=\ell}^n a_{jk} \frac{d_k}{d_j} \\ &= \sum_{k=1}^{\ell-1} (x_k - 1) a_{jk} \frac{d_k}{d_j} + \sum_{k=1}^n a_{jk} \frac{d_k}{d_j} \\ &\leq b \left( \sum_{k=1}^{\ell-1} x_k - (\ell - 1) \right) + M_\ell. \end{aligned} \quad (2.2)$$

Let

$$\phi_\ell = \frac{M_\ell - b + \sqrt{(M_\ell + b)^2 + 4b \sum_{i=1}^{\ell-1} (M_i - M_\ell)}}{2}.$$

For  $1 \leq i \leq \ell - 1$  let

$$x_i = 1 + \frac{M_i - M_\ell}{\phi_\ell + b} \geq 1. \quad (2.3)$$

Then for  $1 \leq i \leq \ell - 1$  we have

$$\begin{aligned} r_i &\leq \frac{b}{x_i} \left( \sum_{k=1, k \neq i}^{\ell-1} x_k - (\ell - 2) \right) + \frac{1}{x_i} M_i \\ &= \frac{b \sum_{k=1}^{\ell-1} (M_k - M_\ell) + \phi_\ell M_i + b M_\ell}{\phi_\ell + b + M_i - M_\ell} \\ &= \frac{\frac{1}{4} [(M_\ell - b)^2 + (M_\ell + b)^2 + 4b \sum_{k=1}^{\ell-1} (M_k - M_\ell) - 2M_\ell^2 - 2b^2 + 4bM_\ell] + \phi_\ell M_i}{\phi_\ell + b + M_i - M_\ell} \end{aligned}$$

$$\begin{aligned} &= \frac{\phi_\ell^2 + \phi_\ell b - \phi_\ell M_\ell + \phi_\ell M_i}{\phi_\ell + b + M_i - M_\ell} \\ &= \phi_\ell. \end{aligned}$$

For  $\ell \leq j \leq n$  we have

$$\begin{aligned} r_i &\leq b \left( \sum_{k=1}^{\ell-1} x_k - (\ell - 1) \right) + M_\ell \\ &= \frac{b \sum_{k=1}^{\ell-1} (M_k - M_\ell) + \phi_\ell M_\ell + b M_\ell}{\phi_\ell + b} \\ &= \frac{\frac{1}{4} [4b \sum_{k=1}^{\ell-1} (M_k - M_\ell) + 2M_\ell \sqrt{(M_\ell + b)^2 + 4b \sum_{k=1}^{\ell-1} (M_k - M_\ell) + 2M_\ell^2 + 2bM_\ell}]}{\phi_\ell + b} \\ &= \frac{\phi_\ell^2 + \phi_\ell b}{\phi_\ell + b} \\ &= \phi_\ell. \end{aligned}$$

Hence by Theorem 1.1,

$$\rho(G) = \rho(B) \leq \max_{1 \leq i \leq n} \{r_i\} \leq \phi_\ell.$$

The first part of Theorem 1.4 follows.

Suppose  $M_1 = M_2 = \dots = M_n$ . Then  $\rho(G) = M_1 = \phi_\ell$  by Theorem 1.2. Hence the equality in Theorem 1.4 follows.

To prove the necessary condition, suppose  $\rho(G) = \phi_\ell$ . Applying Theorem 1.1 and the inequalities in (2.1) and (2.2),  $\phi_\ell = \rho(G) \leq \max_{1 \leq i \leq n} r_i \leq \phi_\ell$ . Hence  $r_1 = r_2 = \dots = r_n = \phi_\ell$ , and the equalities in (2.1) and (2.2) hold. In particular,

$$b = a_{ik} \frac{d_k}{d_i} \tag{2.4}$$

for any  $1 \leq i \leq n$  and  $1 \leq k \leq \ell - 1$  with  $x_k - 1 > 0$ , and  $M_\ell = M_n$ . We consider three cases:

- (1) Suppose  $M_1 = M_\ell$ : Clearly  $M_1 = M_n$ .
- (2)  $M_{t-1} > M_t = M_\ell$  for some  $3 \leq t \leq \ell$ : Then  $x_k > 1$  for  $1 \leq k \leq t - 1$  by (2.3). Hence by (2.4)

$$b = a_{12} \frac{d_2}{d_1} = a_{21} \frac{d_1}{d_2} = 1,$$

and  $d_i = n - 1$  for all  $i = 1, 2, \dots, n$ . This implies that  $G$  is regular, a contradiction.

- (3)  $M_1 > M_2 = M_\ell$ : Then  $x_1 > 1$  by (2.3). Hence by (2.4),  $b = a_{i1} d_1 / d_i$  for  $2 \leq i \leq n$ . Hence  $d_1 = n - 1$  and  $d_2 = d_3 = \dots = d_n = (n - 1) / b$ . Then  $(n - 1) / b = M_1 > M_2 = M_n = (n - 1) / b - 1 + b$ . This implies  $b < 1$ , a contradiction.

This completes the proof of the theorem. □

**Example 2.1.** The graphs in Figs. 1–3 are pseudo-regular but not regular. In particular, the graph in Fig. 3 has a cycle  $C_k$  of  $k$  vertices, and shares each vertex of  $C_k$  with a triangle  $K_3$ .

An interesting problem could be characterizing all the nonregular pseudo-regular graphs.




Fig. 1. Graph with  $M_i = 2$ .




Fig. 2. Graph with  $M_i = 3$ .




Fig. 3. Graphs with  $M_i = 3$ .

**3. The Shape of the Sequence  $\phi_1, \phi_2, \dots, \phi_n$**

Given a decreasing sequence  $M_1 \geq M_2 \geq \dots \geq M_n$  of positive integers, consider the functions

$$\phi_\ell(x) = \frac{M_\ell - x + \sqrt{(M_\ell + x)^2 + 4x \sum_{i=1}^{\ell-1} (M_i - M_\ell)}}{2}$$

for  $x \in [1, \infty)$ . Note that  $\phi_\ell(b)$  is the upper bound of  $\rho(G)$  in Theorem 1.4.

The following proposition shows that the smaller the  $b$  in Theorem 1.4 is, the lower the upper bound of  $\rho(G)$  reaches.

**Proposition 3.1.** *For any  $1 \leq \ell \leq n$ ,  $\phi_\ell(x)$  is increasing on  $[1, \infty)$ .*

**Proof.** For convenience, let

$$S = \sum_{i=1}^{\ell-1} (M_i - M_\ell).$$

To show that  $\phi_\ell(x)$  is increasing on  $[1, \infty)$ , it is sufficient to show that the derivative of  $\phi_\ell(x)$  is nonnegative. This follows from the following equivalent steps.

$$\begin{aligned} \phi'_\ell(x) &\geq 0 \\ \Leftrightarrow -1 + \frac{M_\ell + x + 2S}{\sqrt{(M_\ell + x)^2 + 4Sx}} &\geq 0 \\ \Leftrightarrow \frac{M_\ell + x + 2S}{\sqrt{(M_\ell + x)^2 + 4Sx}} &\geq 1 \\ \Leftrightarrow (M_\ell + x + 2S)^2 &\geq (M_\ell + x)^2 + 4Sx \\ \Leftrightarrow 4SM_\ell + 4S^2 &\geq 0. \quad \square \end{aligned}$$

Note that for  $1 \leq s \leq n - 1$ ,  $M_s = M_{s+1}$  implies  $\phi_s(x) = \phi_{s+1}(x)$ . We adopt the same viewpoint as [9, Proposition 3.1] to describe when the bound gets improved throughout the sequence  $\phi_1(x), \phi_2(x), \dots, \phi_n(x)$  in the following proposition.

**Proposition 3.2.** *Suppose  $M_s > M_{s+1}$  for some  $1 \leq s \leq n - 1$ , and let the symbol  $\succeq$  denote  $>$  or  $=$ . Then*

$$\phi_s(x) \succeq \phi_{s+1}(x) \quad \text{if and only if} \quad \sum_{i=1}^s M_i \succeq xs(s - 1).$$

**Proof.** Consider the following equivalent relations step by step.

$$\begin{aligned} \phi_s(x) &> \phi_{s+1}(x) \\ \Leftrightarrow M_s - M_{s+1} + \sqrt{(M_s + x)^2 + 4x \sum_{i=1}^{s-1} (M_i - M_s)} &> M_{s+1} - M_{s+2} + \sqrt{(M_{s+1} + x)^2 + 4x \sum_{i=1}^s (M_i - M_{s+1})} \end{aligned}$$

$$\begin{aligned}
 &> \sqrt{(M_{s+1} + x)^2 + 4x \sum_{i=1}^s (M_i - M_{s+1})} \\
 \Leftrightarrow &\sqrt{(M_s + x)^2 + 4x \sum_{i=1}^{s-1} (M_i - M_s)} > 2xs - (M_s + x) \\
 \Leftrightarrow &(M_s + x)^2 + 4x \sum_{i=1}^s (M_i - M_s) > 4x^2s^2 - 4xs(M_s + x) + (M_s + x)^2 \\
 \Leftrightarrow &\sum_{i=1}^s M_i > xs(s - 1),
 \end{aligned}$$

where the third relation is obtained from the second by taking square on both sides, simplifying it, and deleting the common term  $M_s - M_{s+1}$ . Note that even if  $2xs - (M_s + x) < 0$  in the third relation, squaring both sides would be proper since then  $\sqrt{(M_s + x)^2 + 4x \sum_{i=1}^{s-1} (M_i - M_s)} \geq |M_s + x| \geq |2xs - (M_s + x)|$ . Similarly, note that if  $\sum_{i=1}^s M_i = xs(s - 1)$ , then  $M_s \leq xs$  and  $2xs - (M_s + x) \geq 0$ . Hence

$$\phi_s(x) = \phi_{s+1}(x) \tag{3.1}$$

$$\begin{aligned}
 \Leftrightarrow &\sqrt{(M_s + x)^2 + 4x \sum_{i=1}^{s-1} (M_i - M_s)} = 2xs - (M_s + x) \\
 \Leftrightarrow &(M_s + x)^2 + 4x \sum_{i=1}^s (M_i - M_s) = 4x^2s^2 - 4xs(M_s + x) + (M_s + x)^2 \\
 \Leftrightarrow &\sum_{i=1}^s M_i = xs(s - 1). \quad \square
 \end{aligned}$$

The following corollary shows that Theorem 1.3 is an improvement of Theorem 1.2.

**Corollary 3.3.** *For any  $x \in [1, \infty)$ ,  $\phi_2(x) \leq M_1$  with equality if and only if  $M_2 = M_1$ .*

**Proof.** If  $M_2 = M_1$  then  $\phi_2(x) = M_2 \leq M_1$ . Suppose  $M_2 < M_1$ . Choose  $s = 1$  and the symbol  $\succeq$  to be  $>$  in Proposition 3.2,

$$M_1 = \phi_1(x) > \phi_2(x). \quad \square$$

Choosing  $b = \max\{d_i/d_j \mid i \sim j\}$ , by Proposition 3.2 with  $s = 2$  and  $x = b$ , if  $M_2 > M_3$  and  $M_1 + M_2 > 2b$ , then  $\phi_2(b) > \phi_3(b)$ . This is a case when Theorem 1.4 is truly an improvement of Theorem 1.3.




Fig. 4. Graph with  $\phi_2 > \phi_3$ .

**Example 3.4.** For the graph in Fig. 4,  $M_1 = M_2 = 4$ ,  $M_3 = 7/2$ ,  $b = 4/3$ ,  $\phi_1(b) = \phi_2(b) = 4$ ,  $\phi_3(b) \doteq 3.762$  and  $\rho(G) = 1 + \sqrt{7} \doteq 3.646$ .

Note that  $\phi_1(x) = M_1 \geq \phi_2(x)$  by Corollary 3.3, and for  $2 \leq t \leq n - 1$ ,  $\sum_{i=1}^t M_i < xt(t - 1)$  implies  $M_t < x(t - 1)$ , and hence  $\sum_{i=1}^{t+1} M_i < xt(t - 1) + x(t - 1) < xt(t + 1)$ . This implies that the sequence  $\phi_1(x), \phi_2(x), \dots, \phi_n(x)$  is composed by two parts. The first part is decreasing and the second part is increasing. In particular, if we choose  $x = M_1$ ,  $M_2 > M_3$ ,  $s = 2$  and  $\succeq$  to be  $>$  in Proposition 3.2, then  $M_1 + M_2 \not\geq 2M_1 = xs(s - 1)$ , so  $\phi_2(M_1) \leq \phi_3(M_1)$ . Hence  $\phi_2(M_1)$  is smallest among  $\phi_1(M_1), \phi_2(M_1), \dots, \phi_n(M_1)$ .

### Acknowledgments

The authors thank the anonymous referees for giving many valuable suggestions in the presentation of the paper. The first author was partially supported by National Science Council under Grant NSC102-2811-M-214-001. The second author was partially supported by National Science Council under Grant NSC102-2115-M-009-009-MY3.

### References

- [1] A. Berman and X. D. Zhang, On the spectral radius of graphs with cut vertices, *J. Combin. Theory Ser. B* **83** (2001) 233–240.
- [2] R. A. Brualdi and A. J. Hoffman, On the spectral radius of  $(0, 1)$ -matrices, *Linear Algebra Appl.* **65** (1985) 133–146.
- [3] Y. H. Chen, R. Y. Pan and X. D. Zhang, Two sharp upper bounds for the signless Laplacian spectral radius of graphs, *Discrete Math. Algorithms Appl.* **3**(2) (2011) 185–191.
- [4] K. C. Das, Proof of conjecture involving the second largest signless Laplacian eigenvalue and the index of graphs, *Linear Algebra Appl.* **435** (2011) 2420–2424.
- [5] Y. Hong, Upper bounds of the spectral radius of graphs in terms of genus, *J. Combin. Theory Ser. B* **74** (1998) 153–159.
- [6] Y. Hong, J. L. Shu and K. F. Fang, A sharp upper bound of the spectral radius of graphs, *J. Combin. Theory Ser. B* **81** (2001) 177–183.

- [7] J. Li, W. C. Shiu, W. H. Chan and A. Chang, On the spectral radius of graphs with connectivity at most  $k$ , *J. Math. Chem.* **46** (2009) 340–346.
- [8] C. Li, H. J. Wang and P. V. Mieghem, Bounds for the spectral radius of a graph when nodes are removed, *Linear Algebra Appl.* **437** (2012) 319–323.
- [9] C.-A. Liu and C.-W. Weng, Spectral radius and degree sequence of a graph, *Linear Algebra Appl.* **438** (2013) 3511–3515.
- [10] H. Minc, *Nonnegative Matrices* (John Wiley and Sons, New York, 1988).
- [11] J. L. Shu and Y. R. Wu, Sharp upper bounds on the spectral radius of graphs, *Linear Algebra Appl.* **377** (2004) 241–248.
- [12] R. P. Stanley, A bound on the spectral radius of graphs with  $e$  edges, *Linear Algebra Appl.* **87** (1987) 267–269.
- [13] A. M. Yu, M. Lu and F. Tian, On the spectral radius of graphs, *Linear Algebra Appl.* **387** (2004) 41–49.